

A Hamiltonian Decomposition of  $K_{2m}^*$ ,  $2m \geq 8$ 

TIMOTHY W. TILLSON

*Hewlett Packard Company, 3404 E. Harmony Rd., Ft. Collins, Colorado 80525**Communicated by Alan Hoffman*

Received May 17, 1977

It is shown that  $K_{2m}^*$ ,  $2m \geq 8$ , can be decomposed into Hamiltonian circuits. A direct construction utilizing difference methods is given for  $2m \equiv 0 \pmod{4}$ . The case  $2m \equiv 2 \pmod{4}$  is handled inductively by means of a construction which shows that  $K_{4m-2}^*$  admits such a decomposition if  $K_{2m}^*$  does.

In this paper a constructive proof utilizing difference methods will be given of the following conjectures due to Bermond and Faber:

*Conjecture 1.* For  $2m \geq 8$ , the edges of the complete directed graph on  $2m - 1$  vertices  $K_{2m-1}^*$  can be partitioned into  $2m - 1$  directed Hamiltonian paths  $P_1, P_2, \dots, P_{2m-1}$ .

*Conjecture 2.* For  $2m \geq 8$ , the edges of the complete directed graph on  $2m$  vertices  $K_{2m}^*$  can be partitioned into  $2m - 1$  directed Hamiltonian circuits  $C_1, C_2, \dots, C_{2m-1}$ .

These conjectures are made in the article of Bermond and Faber [2], "Decomposition of the Complete Directed Graph into  $k$ -Circuits," where they are shown to be equivalent. They note that the conjecture is false for  $2m = 4$  and 6, while the existence of sequenceable groups of orders 21, 39, 55 and 57 give solutions to (2) for  $2m = 22, 40, 56$  and 58. The authors of this article have also verified (2) for  $8 \leq 2m \leq 18$  by computer. Keedwell has obtained a sequencing of the non-abelian group of order 27 in [3], giving a solution for  $2m = 28$ .

The fact that the edges of  $K_{2m+1}^*$ ,  $m \geq 1$ , can be decomposed into Hamiltonian cycles was known to Kirkman [1]. A type (2) decomposition for  $K_{2m+1}^*$ ,  $m \geq 1$ , is an immediate corollary of this result. One such decomposition, which will be used in the proof of Proposition 2, will now be described.

A Latin square  $L$  of order  $n$  is called *row complete* if for all ordered pairs  $(u, v)$  of distinct symbols there exist integers  $r$  and  $c$ ,  $1 \leq r \leq n$ ,  $1 \leq c \leq n - 1$ , such that  $u = L(r, c)$  and  $v = L(r, c + 1)$ . In [4], Williams

published a construction for row complete latin squares of order  $n = 2m$ ,  $m \geq 1$ , on the symbol set  $\{0, 1, \dots, n-1\}$ . It is given by

$$\begin{aligned} L(1, c) &= c/2 \pmod{n}, & \text{if } c \text{ is even} \\ &= ((c-1)/2) \pmod{n}, & \text{if } c \text{ is odd,} \\ L(r, c) &= r-1 + L(1, c) \pmod{n}, & 1 \leq c \leq n, 2 \leq r \leq n. \end{aligned}$$

The verification that  $L$  is a row complete latin square is easy and is omitted. If we regard the symbols  $\{0, 1, \dots, n-1\}$  as vertices and pairs of adjacent symbols in a row as edges, this construction gives a decomposition of type (1) for  $K_{2m}^*$ ,  $m \geq 1$ . By adjoining the new symbol  $\psi$  in column  $n+1$  of each row of a row complete latin square on  $n = 2m$  symbols, and forming cycles out of each row, we have constructed a decomposition of type (2) for  $K_{2m+1}^*$ ,  $m \geq 1$ .

A direct construction of the decomposition of  $K_{2m}^*$  when  $2m \equiv 0 \pmod{4}$  will be given in Proposition 2. The method is suggested by the following construction of a type (1) decomposition for  $2m = 10$ . We start with a type (2) decomposition of  $K_9^*$ , given in Fig. 1 by a row complete latin square of order

$\psi$	0	1	7	2	6	3	5	4
$\psi$	1	2	0	3	7	4	6	5
$\psi$	2	3	1	4	0	5	7	6
$\psi$	3	4	2	5	1	6	0	7
$\psi$	4	5	3	6	2	7	1	0
$\psi$	5	6	4	7	3	0	2	1
$\psi$	6	7	5	0	4	1	3	2
$\psi$	7	0	6	1	5	2	4	3

Rows are regarded as circuits.

FIG. 1. Type (2) decomposition of  $K_9^*$ .

eight with the symbol  $\psi$  adjoined in the first column of every row. As before, the rows are regarded as being cyclically linked. The set  $\alpha$  of underlined edges has the property that each circuit contains exactly one of its elements, and that it is the edge set of a Hamiltonian path in  $K_9^*$ . If we delete the underlined edge from each of the existing circuits the eight resulting Hamiltonian paths together with the path whose edge set is  $\alpha$  form a type (1) decomposition of  $K_9^*$ .

**PROPOSITION 2.** For  $2m \equiv 0 \pmod{4}$ ,  $2m \geq 8$ , Conjecture 1 is true.

*Proof.* In this and the proofs that follow,  $V(G)$  and  $E(G)$  will denote the vertex and edge sets, respectively, of a graph  $G$ .



By the previous discussion, we have a decomposition of type (2) for  $K_{2m-1}^*$ , that is, a partition of the edges into  $2m-2$  directed Hamiltonian circuits  $C_1, C_2, \dots, C_{2m-2}$ . This construction, using a row complete latin square of order  $n = 2m - 2 = 4k + 2$ ,  $k \geq 1$ , is illustrated in Fig. 2.

The proof consists of exhibiting a set of edges  $\alpha = \{e_i | 1 \leq i \leq 2m-2\}$  in this construction (which are underlined in Fig. 2) such that

$$e_i \in E(C_i), \quad 1 \leq i \leq 2m-2 \quad (3)$$

and

$$\alpha \text{ is the edge set of a directed Hamiltonian path in } K_{2m-1}^*. \quad (4)$$

As before, by deleting  $e_i$  from  $C_i$  for each  $i$ , and setting  $\hat{C}_i = C_i - e_i$ , we obtain a partition  $E(\hat{C}_1) \cup E(\hat{C}_2) \cup \dots \cup E(\hat{C}_{2m-2}) \cup \alpha$  of the edges of  $K_{2m-1}^*$  into  $2m-1$  directed Hamiltonian paths.

Inspection of Fig. 2 shows immediately that  $\alpha$  satisfies property (3).

One way to see that (4) holds for  $\alpha$  follows. To any edge  $e$  in  $E(K_{2m-1}^*)$  not incident with the vertex  $\psi$  we can associate its difference  $d(e) = \text{head}(e) - \text{tail}(e) \pmod{n = 4k + 2}$ . Let  $\beta = \{e \in E(K_{2m-1}^*) | d(e) = 2k-1 \pmod{4k+2}\}$ ; these are the edges formed by columns  $2k-1$  and  $2k$  in Fig. 2. Since  $2k-1$  and  $4k+2$  are relatively prime,  $\beta$  is the edge set of a directed  $(4k+2)$ -gon  $R$  in  $K_{2m-1}^*$ , with  $V(R) = V(K_{2m-1}^*) - \{\psi\}$ . The exact form of  $R$  depends upon the congruence class of  $k \pmod{4}$ , and can be determined by solving congruences of the type  $a + x(2k-1) \equiv b \pmod{4k+2}$  where  $x$  gives the number of edges of  $R$  in the path ( $a = a_0, a_1, a_2, \dots, a_{x-1}, a_x = b$ ). These computations are facilitated by the relations given in the following table.

TABLE I  
Useful Relations mod  $4k+2$

Value of $k \pmod{4}$	$(2k-1)^{-1}$	$k^2$
0	$-k-1$	$-(k/2)$
1	$k$	$(k+1)/2$
2	$-k-1$	$-(k/2) + 2k+1$
3	$k$	$(k+1)/2 + 2k+1$

As an example, if  $k \equiv 1 \pmod{4}$  then by solving  $0 + x_1(2k-1) \equiv 3k+3 \pmod{4k+2}$  using the relations given above, we see that there are  $x_1 = (k-1)/2$  edges of  $R$  in the directed path from vertex 0 to vertex  $3k+3$ .

Let  $\beta = \{(3k+3, k), (2, 2k+1), (k+3, 3k+2), (2k+3, 0)\}$  denote the four edges of  $R$  not in  $\alpha$  and set  $\gamma = \alpha \setminus E(R)$ . The two different orders of occurrence of the edges of  $\beta$  in  $R$  as a function of  $k \pmod{4}$  are shown in Fig. 3a and b; the edges in  $\beta$  are dashed.




FIG. 3. (a)  $R$  when  $k \equiv 1$  or  $2 \pmod{4}$ . The four edges of  $R$  not in  $\alpha$  are dashed. (b)  $R$  when  $k \equiv 0$  or  $3 \pmod{4}$ . The four edges of  $R$  not in  $\alpha$  are dashed.

Figure 4 shows  $R$  with the edges in  $\beta$  deleted and the edges in  $\gamma$  added, that is, the simple open directed path  $P$  with edge set  $\alpha$ . It is clear that  $P$  is Hamiltonian since  $|E(P)| = |E(R)| - 4 + 4 = 4k + 2$  and  $V(P) = V(R) \cup \{\psi\} = V(K_{2m-1}^*)$ . Thus (4) holds and the proof of Proposition 2 is complete.

Let  $K_{S, \bar{S}}$  be the complete bipartite undirected graph with vertex set  $S \cup \bar{S}$ , where we take  $\bar{S} = (\bar{x} | x \in S)$ . The cocktail party graph  $C_{S, \bar{S}}$  is defined by  $V(C_{S, \bar{S}}) = V(K_{S, \bar{S}})$  and  $E(C_{S, \bar{S}}) = E(K_{S, \bar{S}}) - M$ , where  $M$  is the edge set of a perfect matching of  $K_{S, \bar{S}}$ , say  $M = \{\{x, \bar{x}\} | x \in S\}$ .  $K_{S, \bar{S}}^*$  and  $C_{S, \bar{S}}^*$  will denote




FIG. 4. (a)  $R$  as in Fig. 3a with the four edges in  $\beta$  deleted and the four edges in  $\gamma$  added. (b)  $R$  as in Fig. 3b with the four edges in  $\beta$  deleted and the four edges in  $\gamma$  added.

the complete directed and directed cocktail party graphs with vertex set  $S \cup \bar{S}$ ; if  $|S| = p$ , we will also use the notation  $K_{p,p}^*$  and  $C_{p,p}^*$  for these graphs.

**PROPOSITION 3.** *For  $m \geq 2$ , the edges of  $C_{2m-1, 2m-1}^*$  can be partitioned into  $2m - 2$  directed Hamiltonian cycles.*

*Proof.* This proposition is a consequence of the well-known result that if  $n \geq 3$  is odd,  $K_{n,n}$  can be decomposed into hamiltonian cycles and a perfect matching.

**PROPOSITION 4.** *If  $K_{2m}^*$  admits decomposition (2), then so does  $K_{4m-2}^*$ .*

*Proof.* Let  $V(K_{4m-2}^*) = S \cup \bar{S}$ , where we take  $S = \{0, 1, \dots, 2m-3, \infty\}$ , with  $\bar{S}$  and  $C_{2m-1, 2m-1}^*$  as above. By Proposition 3 all we need do is partition  $E(K_{4m-2}^*) - E(C_{2m-1, 2m-1}^*)$  into  $2m - 1$  directed Hamiltonian cycles.

By the hypothesis and the equivalence of (1) and (2) we have a decomposition of  $K_{2m-1}^*$  into  $2m - 1$  directed Hamiltonian paths  $P_1, P_2, \dots, P_{2m-1}$ . Set  $V(K_{2m-1}^*) = S$ . We can define  $2m - 1$  directed Hamiltonian paths  $\bar{P}_1, \bar{P}_2, \dots, \bar{P}_{2m-1}$  on the vertex set  $\bar{S}$  by

$$\begin{aligned} V(\bar{P}_i) &= \{\bar{z} : z \in V(P_i)\}, \\ E(\bar{P}_i) &= \{(\bar{x}, \bar{y}) : (y, x) \in E(P_i)\}, \quad 1 \leq i \leq 2m - 1. \end{aligned}$$

We can link  $P_i$  and  $\bar{P}_i$  to form  $2m - 1$  directed Hamiltonian circuits  $H_i$  of  $E(K_{4m-2}^*) - E(C_{2m-1, 2m-1}^*)$  by adding the edges  $(T(P_i), I(\bar{P}_i))$  and  $(T(\bar{P}_i), I(P_i))$ ,  $1 \leq i \leq 2m - 1$ . Note that the added edges are the set  $M^* = \{(x, \bar{x}), (\bar{x}, x) : x \in S\}$ .

It is clear that  $E(H_i) \cap E(H_j) = \emptyset$  for  $i \neq j$ ,  $1 \leq i, j \leq 2m - 1$ , since  $E(P_i) \cap E(P_j) = E(\bar{P}_i) \cap E(\bar{P}_j) = \emptyset$  and  $\{T(P_i) : 1 \leq i \leq 2m - 1\} = \{I(P_i) : 1 \leq i \leq 2m - 1\} = S$ , as was established in the introductory remarks.

**THEOREM.** *For  $2m \geq 8$ , the edges of the complete directed graph on  $2m$  vertices  $K_{2m}^*$  can be partitioned into  $2m - 1$  directed Hamiltonian circuits  $C_1, C_2, \dots, C_{2m-1}$ .*

*Proof.* For  $m = 4$  the theorem is true by Proposition 2. The solution for  $m = 5$  has already been given in the introductory remarks. We now proceed by induction on  $m$ ,  $m \geq 5$ . If  $2m \equiv 0 \pmod{4}$  we are done by Proposition 2. Suppose  $2m \equiv 2 \pmod{4}$ , say  $2m = 4l - 2$  where  $l \geq 4$ . Since  $2l < 2m$ , the theorem is true for  $K_{2l}^*$  by either the remarks above or the inductive hypothesis. Proposition 4 then gives the result for  $K_{4l-2}^* = K_{2m}^*$ .

## ACKNOWLEDGMENTS

The author would like to thank Richard M. Wilson for his advice and encouragement while writing this paper. The Instructional and Research Computing Center at The Ohio State University provided computer time which assisted in this research.

## REFERENCES

1. BERGE, "The Theory of Graphs," p. 187, Methuen, London, 1958.
2. J. C. BERMOND AND V. FABER, Decomposition of the complete directed graph into  $k$ -circuits, *J. Combinatorial Theory B* **21** (1976), 146–155.
3. A. D. KEEDWELL, Some problems concerning complete latin squares, in "Proceedings, British Combinatorial Conference, 1973," London Mathematical Society Lecture Notes No. 13, pp. 89–96, Cambridge University, 1974.
4. E. J. WILLIAMS, Experimental designs balanced for the estimation of residual effects of treatments, *Austral. J. Sci. Res. Ser. A* **2** (1949), 149–168.